

# Journal Pre-proof

A narrative review of intermittent fasting with exercise

Kelsey Gabel, PhD RD, Assistant Professor, Alyshia Hamm, RD, MPH, Research coordinator, Ola Czyzewski, BS, Graduate Research Associate, Julienne Sanchez Perez, MD, T32 Fellow, Endocrinology, Anisa Fought-Boudaia, BS, Graduate Research Associate, Robert W. Motl, PhD, Professor, Paul R. Hibbing, PhD, Assistant Professor



PII: S2212-2672(24)00254-5

DOI: <https://doi.org/10.1016/j.jand.2024.05.015>

Reference: JAND 55931

To appear in: *Journal of the Academy of Nutrition and Dietetics*

Received Date: 15 November 2023

Revised Date: 23 May 2024

Accepted Date: 30 May 2024

Please cite this article as: Gabel K, Hamm A, Czyzewski O, Perez JS, Fought-Boudaia A, Motl RW, Hibbing PR, A narrative review of intermittent fasting with exercise, *Journal of the Academy of Nutrition and Dietetics* (2024), doi: <https://doi.org/10.1016/j.jand.2024.05.015>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright © 2024 by the Academy of Nutrition and Dietetics.

## **A narrative review of intermittent fasting with exercise**

Kelsey Gabel, PhD RD<sup>1,2</sup>, Alyshia Hamm<sup>2</sup>, Ola Czyzewski<sup>1</sup>, Julienne Sanchez Perez<sup>1,3</sup> MD, Anisa Fought-Boudaia<sup>1</sup>, Robert W Motl<sup>1,2</sup> PhD, Paul R. Hibbing<sup>1</sup> PhD

### **Author affiliations:**

<sup>1</sup> Department of Kinesiology and Nutrition, University of Illinois at Chicago, Chicago, IL, USA

<sup>2</sup> University of Illinois Cancer Center, University of Illinois at Chicago, Chicago, IL, USA

<sup>3</sup> Department of Medicine, Endocrinology, University of Illinois at Chicago, Chicago, IL USA

### **Author Positions:**

Kelsey Gabel PhD, RD; Assistant Professor

Alyshia Hamm<sup>2</sup> RD, MPH; Research coordinator,

Ola Czyzewski<sup>1</sup> BS, Graduate Research Associate

Julienne Sanchez Perez<sup>1,3</sup> MD, T32 Fellow, Endocrinology

Anisa Fought-Boudaia<sup>1</sup> BS, Graduate Research Associate

Robert W Motl<sup>1,2</sup> PhD, Professor

Paul R. Hibbing<sup>1</sup> PhD, Assistant Professor

### **Correspondence and reprint requests:**

Kelsey Gabel RD, PhD, Assistant Professor of Nutrition

Department of Kinesiology and Nutrition, University of Illinois at Chicago

1919 West Taylor Street, Room 532, Chicago, IL, 60612

Tel: 312-413-8911, Email: kdipma2@uic.edu

**Author Contributions:** KG collected the data and wrote the first draft with contributions from AH, OC, JSP, AF, RWM, and PRH. All authors reviewed and revised subsequent drafts of the manuscript.

**Funding:** National Institutes of Health, K12HD101373 BIRCWH (KG), Roybal 5P30AG022849 (KG).  
No other disclosures of conflict of interest exist.

Conflict of interest disclosure: The authors have no conflicts of interest to disclose.

**Word count: Abstract:** 293; **Manuscript:** 6,354

**Key words:** intermittent fasting, alternate day fasting, time restricted eating, exercise, physical activity

## 1 **RESEARCH SNAPSHOT**

2 **Research question:** How does the combination of exercise and intermittent fasting affect body  
3 weight, body composition, cardiometabolic risk factors, and physical fitness?

4 **Key Findings:** When combined with different modalities of exercise, intermittent fasting can  
5 reduce body weight and fat mass while eliciting training adaptations. Evidence is equivocal  
6 regarding the impact on lean mass and cardiometabolic markers, and there is a need for longer  
7 and better-powered interventions in this area. Combining intermittent fasting with exercise may  
8 provide an accessible, low burden alternative to traditional caloric restriction. Future trials should  
9 prioritize recruitment of well-powered samples comprising both males and females, a broad  
10 range of ages, and those at risk for cardiometabolic disease.

## 11 **ABSTRACT**

12 Intermittent fasting is a dietary pattern that encompasses the 5:2 diet, alternate day fasting  
13 (ADF), and time restricted eating (TRE). All three involve alternating periods of fasting and ad  
14 libitum eating. Like other dietary strategies, intermittent fasting typically induces loss of both fat  
15 mass and lean mass. Exercise may thus be a useful adjuvant to promote lean mass retention  
16 while adding cardiometabolic, cognitive, mental, and emotional health improvements. In this  
17 narrative review, we summarize current evidence regarding the combination of intermittent  
18 fasting and exercise and its impacts on body weight, body composition, cardiometabolic risk, and  
19 muscular and cardiorespiratory fitness. A PubMed search was conducted to identify all trials  
20 lasting >4 weeks that combined 5:2, ADF, or TRE with any modality exercise and had body  
21 weight as an endpoint. A total of 23 trials (26 publications) were identified. Evidence suggests  
22 that combining intermittent fasting with exercise leads to decreased fat mass regardless of weight

23 status. However, evidence is equivocal for the impact on other aspects of weight loss and body  
24 composition, fat free mass and cardiometabolic risk factors and may be dependent on weight  
25 status or exercise dosages (i.e., frequency, intensity, duration, and modality). Higher-powered  
26 trials are needed to determine the efficacy of combining exercise and intermittent fasting for  
27 benefits on bodyweight and cardiometabolic risk. Current evidence suggests that intermittent  
28 fasting does not impair adaptation to exercise training, and may improve explosive strength,  
29 endurance, and cardiopulmonary measures such as maximal oxygen consumption. Additionally,  
30 we discuss limitations in the current evidence base, and opportunities for continued investigation.  
31 Future trials in this area should consider interventions that have 1) increase sample size, 2)  
32 longer intervention duration, 3) broadened inclusion criteria, 4) objective measures of diet and  
33 exercise adherence, and 5) diversity of sample population.

## 34 **INTRODUCTION**

35 Obesity and overweight continues to be a significant issue in the United States (U.S.) with 30%  
36 of Americans having overweight, 40% having obesity, and almost 10% having severe obesity.<sup>1</sup>  
37 While the use of anti-obesity drugs has increased, dietary behavioral interventions remain  
38 necessary for overall health, increased efficacy of pharmacological treatment, and weight loss  
39 maintenance. Intermittent fasting, a dietary behavioral intervention for weight management, has  
40 increased in popularity in the last two decades due to the ease of implementation and less  
41 stringent food restriction requirements than traditional caloric restriction (CR). Three main forms  
42 of intermittent fasting have emerged, namely, the 5:2 diet, alternate day fasting (ADF), and time-  
43 restricted eating (TRE). Each form alternates “feast” periods (ad libitum intake) with “fast”  
44 periods (calorie abstention), yet in different ways. The 5:2 diet consists of 2 consecutive or non-  
45 consecutive “fast days” of 0-500kcal (or up to 25% of energy needs) each week with the

46 remaining five days as “feast days”. Similarly, ADF consists of eating 0-500kcal (or up to 25%  
47 of energy needs) on the “fast day” but this is alternated every other day throughout the week with  
48 ad libitum eating on the “feast day”. TRE consists of ad libitum eating during a 6- to 10-hour  
49 daily window, while fasting the remaining 14-18 hours. Prior research suggests that the 5:2 diet  
50 and ADF may be as efficacious as traditional CR for weight loss,<sup>2,3</sup> with 6-12 weeks of either  
51 approach resulting in weight loss of 3-8% and decreases in blood pressure, insulin resistance, and  
52 other cardiometabolic markers.<sup>4</sup> Evidence is less conclusive for TRE, but 8-12 weeks appears to  
53 result in a calorie deficit of 20-40% and weight loss of 2-4% with uncertain effects on  
54 cardiometabolic outcomes.<sup>5</sup>

55  
56 Recently, interest has grown in the combined effects of intermittent fasting and exercise.  
57 Exercise, planned physical activity, is a key determinant of energy balance and may therefore  
58 result in an augmented effect on body weight while also contributing to improvements in serum  
59 lipids, blood pressure, fasting glucose. The current Physical Activity Guidelines for Americans  
60 call for 150-300 minutes of moderate-intensity aerobic activity per week and two days of muscle  
61 strengthening activities weekly for disease prevention, health promotion, and weight loss.  
62 However, 55% of Americans don’t meet these recommendations,<sup>6,7</sup> which compounds the  
63 cardiovascular and cardiometabolic impact of overweight and obesity. Furthermore, as exercise  
64 is a key regulator of lean mass, the addition of exercise to a dietary regimens such as intermittent  
65 fasting may help to mitigate lean mass loss typically experienced when undertaking energy  
66 restricted diets.<sup>8,9</sup> This is critical to pursue because lean mass (especially skeletal muscle) plays a  
67 central role in regulating basal metabolism and peripheral glucose uptake.<sup>6,10,11</sup> These benefits  
68 underscore the potential benefits of combining exercise with intermittent fasting, but to date

69 there has been limited synthesis of evidence from studies examining the joint effects of  
70 intermittent fasting and exercise on weight loss and related health markers.

71

72 The purpose of this narrative review is to summarize the current literature examining the effects  
73 of intermittent fasting (5:2, ADF, and TRE) combined with various modalities of exercise on  
74 body weight and body composition, cardiometabolic risk, glucoregulatory factors, and muscular  
75 and cardiorespiratory fitness.

## 76 **METHODS**

77 This is a narrative review and as such was not registered. A PubMed search was conducted using  
78 the following key words or MeSH terms: "humans", "fasting", "time restricted eating", "alternate  
79 day fasting", "alternate day modified fasting", "intermittent fasting", "fasting", "intermittent  
80 energy restriction", "exercise", "exercise therapy", "resistance training", "resistance exercise",  
81 "strength training", "aerobic exercise", "exercise", "aerobic training", "physical activity",  
82 "endurance exercise", "weightlifting", "walking". The inclusion criteria for research articles  
83 were as follows: (1) randomized controlled trials and nonrandomized trials; (2) adult male and  
84 female participants (>18 years); (3) endpoints that included changes in body weight; and (4) only  
85 studies that included intermittent fasting and exercise combined. The following exclusion criteria  
86 were applied: (1) cohort, cross sectional, and observational studies; (2) fasting performed as a  
87 religious practice (e.g., Ramadan or Seventh Day Adventist); and (3) trial durations <4 weeks.

## 88 **RESULTS**

89 It is important to consider the effects of intermittent fasting on body weight and composition  
90 since weight loss of  $\geq 5\%$  can reduce cardiometabolic risk in individuals with overweight and

91 obesity.<sup>12</sup> Even in individuals without overweight or obesity a caloric deficit combined with  
92 exercise can improve longevity, cognition, and physical functioning with age.<sup>13,14</sup> Accordingly,  
93 several trials have examined the effects of combining intermittent fasting and various forms of  
94 exercise on body weight and composition in individuals with and without obesity. Our search  
95 retrieved 7 trials (8 publications)<sup>15-22</sup> on 5:2 combined with exercise, 3 trials (4 publications)<sup>23-26</sup>  
96 on ADF combined with exercise, and 13 trials (14 publications) on TRE combined with  
97 exercise.<sup>27-39</sup> **Table 1** describes the trial design and intervention characteristics. **Table 2** describes  
98 the findings on body weight, body composition cardiometabolic factors, and muscular and  
99 cardiorespiratory fitness. **Table 3** describes the heterogeneity between trial participants and  
100 adherence monitoring.

## 101 5:2

102 Seven trials (8 publications) have examined the effect of the 5:2 diet combined with  
103 exercise.<sup>15-22</sup> One was a randomized control trial,<sup>18</sup> 5 trials (6 publications) were randomized  
104 trials without a control,<sup>15-17,19,21,22</sup> and 1 trial was not randomized.<sup>20</sup> One trial included  
105 participants with normal weight, overweight, or obesity,<sup>15,22</sup> five trials included participants with  
106 overweight or obesity,<sup>17-21</sup> and one trial included participants with obesity only.<sup>16</sup> Exercise  
107 interventions included aerobic activity, steps/day, resistance training, high intensity interval  
108 training or a combination of modalities.

109 Batitucci et al.<sup>16</sup> conducted a parallel arm trial examining the 5:2 diet (600kcal fast days),  
110 high intensity interval training (HIIT) three days a week, or the 5:2 diet and HIIT combined.  
111 Only the combination group decreased body weight significantly (-2%) after the 8-week  
112 intervention. Fat mass and waist circumference decreased, and fat free mass increased in both the  
113 combination and HIIT groups but remained unchanged in the 5:2 alone group. A group difference



114 was reported between the two exercise groups compared to diet alone at week 8, however, no  
115 time by diet effect was reported. Kang et al.<sup>17</sup> compared 5:2 (30% of maintenance calories on  
116 fast days), CR alone (30% calorie deficit) and CR plus protein meal replacement (30% calorie  
117 deficit) in a parallel arm randomized trial. All participants were instructed to increase their  
118 physical activity to 150-300 minutes weekly. The 5:2 diet and the meal replacement groups lost  
119 9% of body weight after 12 weeks, which was significantly more weight loss over time  
120 compared to the CR group (-5%, time by diet interaction). Importantly, significantly more  
121 participants lost clinically significant (5% and 10%) body weight from baseline in the 5:2 and  
122 meal replacement groups than the CR group. Fat mass decreased and fat free mass increased in  
123 all three groups with no difference between groups. Hottenrott et al.<sup>18</sup> compared the 5:2 diet to an  
124 unrestricted diet group combined with 30–60 min of running and 20 min of resistance training 3-  
125 4 days a week throughout the 12-week intervention in healthy individuals with obesity.  
126 Additionally, both groups were divided and randomized to ingest an alkaline supplement or  
127 placebo. The 5:2 groups lost significantly more body weight and fat mass than the ad libitum  
128 groups with or without the alkaline supplement over time (time by diet interaction). Additionally,  
129 the 5:2 group combined with the alkaline supplement lost significantly more body weight, fat  
130 mass, and visceral fat mass than the 5:2 diet group alone. Keenan et al.<sup>15</sup> compared 5:2 (30% of  
131 maintenance calories on fast days) or CR (20% calorie deficit) combined with 2 supervised  
132 resistance training sessions and 1 unsupervised aerobic or resistance training session in  
133 individuals with normal weight, overweight, or obesity. Body weight (5:2: -5% males, -3%  
134 females; CR -7% males, -3% females), and fat mass decreased significantly, and fat free mass  
135 significantly increased in both IF and CR groups after 12 weeks. No time by diet interaction was  
136 reported for body weight or composition; however, time by sex interactions were reported with

137 males losing more weight than females and females gaining more lean mass than males. The  
138 increase in lean mass in the females may account for the difference in weight loss between the  
139 sexes. Additionally, the CR group increased muscle surface area significantly more than the 5:2  
140 group over time (time by diet interaction). Cooke et al.<sup>21</sup> examined 5:2, sprint interval training  
141 sessions 3 times a week, or a combination of 5:2 and sprint interval training for 16 weeks in  
142 individuals with overweight or obesity. Body weight and fat mass decreased significantly more  
143 over time in the 5:2 alone or combination group (data not provided) than sprint training alone  
144 (time by diet interaction). At 8 weeks both 5:2 groups lost lean body mass compared to the sprint  
145 group alone, however this was not significant at week 16. Waist circumference decreased in all  
146 three groups after 16 weeks. Headland et al.<sup>19</sup> compared the 5:2 diet (500kcal for females and  
147 600kcal for males on fast days) to a week-on week-off diet (1000kcal/d for females and  
148 1200kcal/d for males) or CR (30% calorie deficit) in individuals with overweight and obesity.  
149 Participants were advised to increase their steps to 10,000 per day. After 52 weeks, all groups  
150 decreased lean mass and fat mass from baseline, resulting in 6-7% mean body weight reduction  
151 in each group with no time by diet interaction. Jospe et al.<sup>20</sup> compared the 5:2 diet, a  
152 Mediterranean diet (mostly plant based foods with mono and polyunsaturated fats), and  
153 paleolithic diet (restriction of grains, legumes, and dairy) in individuals with overweight or  
154 obesity in a non-randomized parallel-arm trial. Participants were able to choose their diet arm  
155 and one of two exercise interventions, 1) 150-300 minutes of aerobic activity plus two days of  
156 resistance training (58% of participants) or 2) at home-based HIIT program (42% of  
157 participants). All three diet groups reported significant weight loss after 24 weeks. However,  
158 only the 5:2 diet and Mediterranean diet groups significantly decreased body weight, fat mass,  
159 and visceral fat mass after 52 weeks of the intervention. Lean mass was not reported in this trial,

160 but when considering changes in body fat percentage and mean changes in weight, it appears that  
161 both the Mediterranean diet and paleolithic diet retained or gained lean mass compared to a loss  
162 in lean mass in the 5:2 group (non-significant). However, these results may be skewed as the 5:2  
163 diet group had more than double the participants of the Mediterranean diet and paleolithic diet  
164 groups.

165

166       Regarding cardiovascular disease risk, four trials have examined the effect of the 5:2 diet  
167 with exercise on fasting lipids, glucose, insulin, and measures of insulin sensitivity or resistance.  
168 <sup>15,19-21</sup> Two of the trials also examined the effect on blood pressure.<sup>20,21</sup> Keenan et al.<sup>15</sup> reported a  
169 significant decrease in LDL cholesterol and HDL cholesterol from baseline in both the 5:2 and  
170 CR groups when combined with resistance training. The 5:2 group decreased LDL cholesterol  
171 significantly more than the CR group over time (time by diet interaction). Additionally, females  
172 decreased HDL significantly more over time than males (time x sex interaction). No changes  
173 were reported for triglycerides, fasting glucose, fasting insulin or insulin resistance via  
174 homeostatic model assessment – insulin resistance ( $HOMA-IR = \frac{\text{fasting plasma insulin}}{\text{fasting plasma glucose}/22.5}$ ).<sup>40</sup> Cooke et al.<sup>21</sup> reported a significant decrease in LDL cholesterol over  
175 time when combining the 5:2 diet with sprint interval training, whereas no changes were  
176 observed in the 5:2 diet alone and sprint interval training alone. HDL cholesterol, triglycerides,  
177 blood pressure, fasting insulin, fasting glucose, and HOMA-IR also remained unchanged in all  
178 three groups. Headland et al.<sup>19</sup> reported a significant increase in HDL cholesterol and a  
179 significant decrease in triglycerides after 52 weeks of 5:2, week on week off, and CR combined  
180 with 10,000 steps/day, with no between group differences reported. Fasting glucose remained  
181 unchanged. Jospe et al.<sup>20</sup> reported a significant decrease in LDL cholesterol in the  
182

183 Mediterranean and paleolithic diets combined with exercise, and an increase in HDL cholesterol  
184 with 5:2 combined with exercise. Triglycerides remained unchanged in all groups. Systolic and  
185 diastolic blood pressure decreased significantly in the Mediterranean and 5:2 diet, whereas only  
186 diastolic blood pressure decreased in the Paleolithic diet group after 52 weeks. The  
187 Mediterranean diet group significantly lowered their HbA1c over time and this changed resulted  
188 in a time by diet interaction.

189  
190 In regards to muscular strength or cardiorespiratory performance, four studies have  
191 examined the effects of 5:2 combined with exercise.<sup>16,18,21,22</sup> Batitucci et al.<sup>16</sup> reported that HIIT  
192 alone and 5:2 combined with HIIT improved shuttle walking test, strength (abdominal test, push  
193 up test, squat test, 1 repetition maximum leg 45° test, 1 repetition maximum bench press test,  
194 dorsal dynamometer, handgrip), observed maximal heart rate, and  $VO_{2max}$  after 8 weeks. Keenan  
195 et al.<sup>22</sup> reported increased upper and lower body strength (3-repetition maximum and volume test  
196 of bench press and leg press) from baseline when combining resistance training with 5:2 or CR.  
197 The CR group increased muscle surface area significantly more than the 5:2 group over time  
198 (time by diet interaction). Hottenrott et al.<sup>18</sup> reported an increase in maximum running velocity  
199 from baseline in the 5:2 group combined with exercise and an alkaline supplement compared to  
200 5:2 combined with exercise and a placebo. No changes were reported in 5:2 alone or the ad  
201 libitum groups independent of the alkaline supplement. Lastly, Cooke et al.<sup>21</sup> reported a  
202 significant increase in  $VO_{2peak}$  with sprint interval training alone and in the combination group  
203 compared with 5:2 alone. The combination group increased  $VO_{2peak}$  significantly more over time  
204 than sprint interval training alone (time by diet interaction).

205 Summary of findings: Altogether, the results of the above studies suggest that the 5:2 diet  
206 combined with exercise appears to produce weight loss of 2-9% and significant decreases in fat  
207 mass after 8-52 weeks of the diet these reductions. Additionally, 5:2 appears to produce similar  
208 weight and fat mass loss to traditional CR or the Mediterranean diet. It is unclear if the 5:2 diet  
209 ameliorates lean mass loss, which may be dependent on the magnitude of caloric deficit and or  
210 modality of exercise. However, it does appear that 5:2 combined with exercise improves lipid  
211 profile, blood pressure, and insulin sensitivity, although the data are not entirely conclusive.  
212 According to the data presented here, diets that affect diet quality, such as the Mediterranean  
213 diet, may be more beneficial for glucose regulation and insulin sensitivity. These data suggest  
214 that training adaptations are not impaired by the caloric restriction of 5:2 and may improve both  
215 strength, running velocity and  $V_{O_{2peak}}$ .

## 216 **ADF**

217 Three trials (4 publications) have examined the effect of ADF combined with exercise on  
218 body weight and body composition.<sup>23-25</sup> All three trials utilized a randomized controlled factorial  
219 design.<sup>23-25</sup> One trial included participants with overweight and obesity,<sup>25,26</sup> whereas the others  
220 included only participants with obesity.<sup>23,24</sup> One trial examined resistance and aerobic  
221 training<sup>25,26</sup> and the other two examined aerobic activity only.<sup>23,24</sup> Cho et al.<sup>25,26</sup> compared ADF,  
222 exercise (resistance and aerobic exercise 3 times a week), and ADF combined with exercise to a  
223 control group. However, this search revealed two manuscripts for NCT03652532, with varying  
224 results.<sup>25,26</sup> The manuscript from Cho et al<sup>25</sup>, analyzed 31 completers. The ADF, exercise, and  
225 combination groups all reduced body weight, body fat percentage, and fat mass significantly  
226 after 8 weeks of the intervention. A significant time by diet interaction was reported for both  
227 ADF and combination groups compared to controls for body weight change and fat mass loss.

228 Skeletal muscle decreased in the combination group from baseline. Under the same  
229 clinicaltrials.gov registration, Oh et al.<sup>26</sup> analyzed 35 completers. Body weight decreased in the  
230 ADF and combination groups. Fat mass and fat free mass decreased in the combination group  
231 only. Waist circumference decreased significantly from baseline in ADF, combination, and  
232 exercise groups. Body fat percentage decreased over time in the ADF group and combination  
233 group, with a significant time by diet interaction between the combination and control groups.  
234 Bhutani et al.<sup>23</sup> compared ADF, aerobic exercise 3 days a week, and ADF and exercise combined  
235 compared to a no intervention control group in individuals with obesity. All three intervention  
236 groups lost a significant amount of body weight from baseline (ADF + exercise: -7%, ADF: -3%,  
237 Exercise: -1%) and decreased waist circumference after 12 weeks. Participants in the  
238 combination group lost significantly more body weight and decreased waist circumference  
239 significantly more than the other groups over time (time by diet interaction). Fat mass decreased  
240 in the ADF and combination group whereas fat-free mass decreased in the ADF group only.  
241 Ezpeleta et al.<sup>24</sup> performed a similar factorial trial to Bhutani et al.,<sup>23</sup> however the exercise dose  
242 was higher (aerobic exercise 5 days/week). Participants had obesity and non-alcoholic fatty liver  
243 disease. After 12 weeks, body weight decreased significantly by -5% in both the ADF and  
244 combination groups, and -2% for exercise alone. Body weight, fat mass, fat free mass, and  
245 visceral fat decreased significantly more in the combination group over time (time by diet  
246 interaction) compared to exercise alone and the controls. However, no differences were reported  
247 between ADF alone or ADF combined with exercise for body weight, fat mass, lean mass, or  
248 visceral fat mass.  
249

250           Regarding cardiometabolic risk, three trials have examined the effect of ADF combined  
251 with exercise on fasting lipids and glucoregulatory factors,<sup>23-25</sup> two of which also examined  
252 effects on blood pressure.<sup>23,24</sup> Cho et al.<sup>25</sup> reported no changes in LDL cholesterol or HDL  
253 cholesterol after 8 weeks of ADF or exercise alone or combined. However, the combination  
254 group significantly decreased triglycerides from baseline whereas the control group increased  
255 triglycerides significantly more over time when compared with the ADF and combination groups  
256 (time by diet interaction). No changes were reported in fasting glucose, fasting insulin, or insulin  
257 resistance. Bhutani et al.<sup>23</sup> reported a significant decrease in LDL cholesterol and an increase in  
258 HDL cholesterol (time by diet interaction compared to controls) in the ADF combined with  
259 exercise groups; ADF and exercise alone remained unchanged. Triglycerides remained  
260 unchanged in all groups. Only the ADF group reported a significant decrease in both systolic  
261 and diastolic blood pressure from baseline however, the ADF group had significantly higher  
262 blood pressure at baseline. Fasting glucose decreased significantly from baseline in all treatment  
263 groups and remained unchanged in the control group. Fasting insulin decreased from baseline in  
264 the ADF group only, while insulin resistance remained unchanged in all groups. Lastly, Ezpeleta  
265 et al.<sup>24</sup> reported no changes in LDL and HDL cholesterol in ADF, exercise alone, or the  
266 combination groups. ADF alone reduced triglycerides significantly from baseline. Diastolic  
267 blood pressure decreased from baseline in the combination group only. Fasting insulin  
268 significantly decreased and insulin sensitivity via the quantitative insulin sensitivity check index  
269 ( $QUICKI = 1 / [\log [\text{insulin (mIU/ml)}] + \log [\text{glucose (mg/dl)}]]^{41}$ ) statistically increased in the  
270 ADF, aerobic exercise, and combination groups, but not the control group. The combination  
271 group also decreased fasting insulin and increased insulin sensitivity significantly more over time  
272 compared to the exercise and control groups (time by diet interaction). No difference was

273 reported between the ADF and combination groups, and HbA1c remained unchanged in all  
274 groups.

275

276         Regarding muscular strength or cardiorespiratory performance only one study has  
277 examined the effect of ADF with exercise on muscular strength or cardiorespiratory  
278 performance. Cho et al.<sup>25</sup> reported a significant increase  $VO_{2max}$  from baseline in the ADF  
279 combined with exercise group. Muscle strength (chest press and pulldown) was significantly  
280 increased in the exercise and combination group from baseline. Chest press significantly  
281 decreased in the ADF alone group while no changes were reported in the controls.

282

283         Summary of findings: These data suggest that ADF combined with aerobic training may  
284 improve body weight by 4-7% in 8-12 weeks as well as significant decreases in fat mass, and  
285 waist circumference. When considering lean mass, the results are incongruent. Cho et al.<sup>25,26</sup> and  
286 Ezpeleta et al.<sup>24</sup> did not report lean mass change with the addition of exercise, however, Bhutani  
287 et al.<sup>23</sup> only reported lean mass loss in the diet alone group indicating that exercise mitigated this  
288 loss. Due to the paucity of data and inconsistency of results, the effect of ADF combined with  
289 exercise is unclear for triglycerides and blood pressure. Regarding glucose regulation, data from  
290 Ezpeleta et al.<sup>24</sup> appears promising, yet it is also uncertain if combining ADF and exercise results  
291 in favorable impact on glucoregulatory factors. Lastly, only one trial examined muscular strength  
292 or cardiorespiratory in ADF combined with exercise. More studies will need to examine if these  
293 data can be replicated.

294



295 **TRE**

296 Thirteen studies (14 publications) have examined the effect of TRE combined with different  
297 modalities of exercise on body weight and body composition.<sup>27-39</sup> Nine<sup>27,29,31,33-37,42</sup> were  
298 randomized control trials wherein controls were prescribed a 12-h eating window or instructed to  
299 maintain current eating patterns, one<sup>38</sup> was a randomized trial with no controls, two were  
300 randomized crossover design<sup>28,39</sup> and one (two publications) was a single arm design.<sup>30,43</sup> Three  
301 trials included participants with overweight or obesity<sup>33,37,38</sup> with the remaining ten trials  
302 including participants with normal weight only.<sup>27-32,34-36,39</sup> Exercise interventions included  
303 aerobic and endurance activity, resistance training, high intensity interval training or a  
304 combination of modalities. Haganes et al.<sup>33</sup> examined TRE, HIIT (3 days per week), or TRE  
305 combined with HIIT compared to controls in individuals with overweight or obesity. Compared  
306 to the control group, the TRE, HIIT, and combination groups significantly reduced their body  
307 weight (TRE: -2%, HIIT: -2%, combination -4%), fat mass, and visceral fat mass from baseline  
308 (time by diet interaction). Fat free mass decreased significantly in the TRE group over time  
309 compared to the control group (time by diet interaction). Isenmann et al.<sup>38</sup> examined 8-h ad  
310 libitum TRE with macronutrient recommendations (45-65% carbohydrate, 20-35% fat, 20-35%  
311 protein) compared to a Macronutrient based diet which consisted of 80% unprocessed foods, in  
312 participants with overweight or obesity for 14 weeks (8 week intervention period and 6 week  
313 independent period). A 500kcal deficit was included during the independent period of the  
314 macronutrient diet. Both groups were asked to follow their diet and attend two training sessions a  
315 week. After 14 weeks, both groups significantly decreased body weight (-5%), fat mass, and  
316 waist circumference from baseline with no differences between groups. No changes were  
317 reported in fat free mass. Kotarsky et al.<sup>37</sup> examined 8-h TRE or a “normal” eating window

318 combined with 300 minutes of moderate or 150 minutes of vigorous aerobic activity and  
319 resistance training on three non-consecutive days per week in individuals with overweight or  
320 obesity. After 8 weeks, the TRE group lost significantly more body weight (-4%) and fat mass  
321 over time compared to the normal eating group (time by diet interaction). Both the TRE and  
322 normal eating group increased fat free mass and decreased waist circumference from baseline  
323 with no differences between groups. Morro et al.<sup>27</sup> compared 8-h TRE compared to a control (12-  
324 h) diet in young healthy male elite cyclists for four weeks. Both groups combined their dietary  
325 intervention with cycling (500km/week in six training sessions/week) and were given a weight  
326 maintenance calorie goal to control for energy intake. The TRE group significantly reduced body  
327 weight (-2%). Fat mass was significantly lower in the TRE group than the controls at week 4  
328 (group difference), however this change was not significant over time or time by diet. No  
329 changes were reported in fat free mass. Richardson et al.<sup>28</sup> also compared isocaloric 8-h TRE to a  
330 12-hour control diet group in healthy male endurance trained runners. Participants were asked to  
331 maintain their current training regimen for both arms of the study. No changes were reported in  
332 body weight or fat free mass after 4 weeks, but fat mass decreased significantly during the TRE  
333 intervention over time. Correia et al.<sup>29</sup> randomized healthy trained young males to TRE or  
334 normal diet (12-h), both of which were combined with 3 resistance training sessions per week.  
335 After 4 weeks, no changes in body weight, fat free mass, or skeletal muscle were reported in  
336 either group. Fat mass decreased in both groups from baseline, with no differences between  
337 groups. Waldman et al.<sup>30,43</sup> examined 8-h TRE in middle-aged competitive male cyclists for 8  
338 weeks. Participants self-selected their eating window to fit their family's eating schedule. All  
339 cyclists reported exercise over 150 min/week and were asked to continue their current habitual  
340 exercise during the dietary intervention. Participants significantly reduced body weight (-3%)

341 and fat mass after 8 weeks. Fat free mass and abdominal skin fold remained unchanged. Morro et  
342 al.<sup>31</sup> compared a modified form of TRE (4-h window on 4 days per week) against a normal diet  
343 (12-h) control in lean healthy young males. Participants were instructed to eat calories for weight  
344 maintenance and perform resistance training on 3 non-consecutive days/week. No changes were  
345 reported in body weight or fat free mass after 8 weeks, but fat mass decreased significantly more  
346 over time in the TRE compared to the normal diet group (time by diet interaction). Moro et al.  
347 then performed a follow-up at one year from baseline (10 months after the completion of the  
348 previous trial). The TRE group significantly decreased their body weight (-3%) and fat mass,  
349 while the normal diet group significantly increased their body weight and fat mass. These  
350 changes were significant over time and reporting a significant time by diet interaction. No  
351 changes were reported in visceral fat mass at follow-up.<sup>42</sup> However, at 12 months, TRE  
352 observed a significant decrease in arm and thigh circumference (cross-sectional area) from  
353 baseline compared to the normal diet group (time by diet interaction). Tinsley et al.<sup>34</sup> recruited  
354 healthy trained males and compared an isocaloric 7-h TRE diet to a normal diet (12-h) control  
355 for 8 weeks. Both groups performed resistance training on three non-consecutive days each week  
356 throughout the trial. Body weight, fat mass, fat free mass, and visceral fat mass remained  
357 unchanged in both groups. Brady et al.<sup>36</sup> examined 8-h TRE versus a control group in 17 male  
358 middle- and long-distance runners who were asked to maintain their habitual exercise. The TRE  
359 group lost significantly more body weight (-3%) than the control at 8 weeks (time by diet  
360 interaction). No changes were reported in fat mass or fat free mass. Tinsley et al.<sup>35</sup> also examined  
361 8-h TRE or normal (12-h) control diet combined with resistance training (three non-consecutive  
362 days per week) in 40 resistance trained females for 8 weeks. Both diet groups were tested with  
363 and without Hydroxymethylbuterate (HMB) supplementation, which may promote muscle

364 growth. All groups increased body weight (1-2%) and fat free mass significantly. Fat mass was  
365 reduced in the TRE groups independent of HMB supplementation. No differences in body  
366 weight, fat mass or fat free mass were reported between groups in the intention to treat analysis.  
367 However, the per protocol analysis (n=24) reported a significant time by diet interaction with  
368 larger reductions in fat mass and body fat percentage in the TRE plus HMB group and significant  
369 increases in fat free mass in all groups. Martinez-Rodriguez et al.<sup>39</sup> examined HIIT (3 times per  
370 week) alone compared to HIIT combined with every other day TRE (<14-h eating window with  
371 first meal close to waking) using a randomized crossover design in 14 active, normal weight  
372 females. HIIT alone had no effect on body weight and body fat, but HIIT combined with TRE  
373 produced a significant reduction (time by diet interaction) in fat mass. Fat free mass remained  
374 unchanged.

375  
376 In regards to cardiovascular risk, eight studies have examined the effect of TRE  
377 combined with exercise on fasting lipids,<sup>28,30,31,33,35,37,42</sup> four on blood pressure,<sup>28,33,35,37</sup> and ten  
378 on glucoregulatory factors.<sup>27,28,30,31,33-37,42</sup> Richardson et al.<sup>28</sup> reported no changes in LDL  
379 cholesterol, HDL cholesterol, triglycerides or blood pressure after 4 weeks of TRE or a normal  
380 diet in male elite endurance runners. Fasting glucose, fasting insulin, insulin resistance (HOMA-  
381 IR), and insulin sensitivity (QUICKI) also remained unchanged. Tinsley et al.<sup>35</sup> reported no  
382 changes in LDL cholesterol, HDL cholesterol, triglycerides, fasting insulin, or fasting glucose  
383 after 8 weeks of TRE or a control diet combined with resistance independent of HMB  
384 supplementation in trained lean females. Diastolic blood pressure significantly decreased in both  
385 the TRE and normal diet groups independent of HMB supplementation. Haganes et al.<sup>33</sup> reported  
386 no changes in LDL cholesterol, triglycerides, blood pressure, fasting glucose, fasting insulin, and

387 insulin resistance (HOMA-IR) after 7 weeks of TRE or HIIT alone or combined in individuals  
388 with overweight or obesity. The combination group reported a greater reduction in HDL  
389 cholesterol over time than the other groups (time by diet interaction). Nocturnal glucose  
390 decreased significantly in the TRE and combination groups and HbA1c decreased significantly  
391 in the combination group compared to controls (time by diet interaction). Waldman et al.<sup>30</sup>  
392 reported no changes in LDL cholesterol or triglycerides, however HDL cholesterol increased  
393 after 4 weeks of 8-h TRE in male cyclists. Fasting glucose significantly decreased while fasting  
394 insulin and insulin resistance (HOMA-IR) remained unchanged. Kotarsky et al.<sup>37</sup> reported no  
395 changes in HDL, blood pressure, fasting insulin or HbA1c after TRE combined with 150-300  
396 minutes of exercise for 8 weeks. Brady et al.<sup>36</sup> also reported no change in triglycerides, fasting  
397 glucose, fasting insulin or insulin resistance (HOMA-IR). Moro et al.<sup>42</sup> reported a significant  
398 increase in HDL cholesterol and a significant reduction in triglycerides, glucose, insulin, and  
399 insulin resistance (HOMA-IR) after 8 weeks of 4h TRE four days per week combined with  
400 resistance training in lean trained males. At a year follow-up of the same participants  
401 significantly decreased LDL cholesterol and increased HDL. Compared to the controls over time,  
402 triglycerides decreased significantly (time by diet interaction). Additionally, fasting glucose,  
403 fasting insulin, and insulin resistance decreased significantly more over time compared to the  
404 normal diet controls (time by diet interaction). In a different trial, Morro et al.<sup>27</sup> reported no  
405 differences in triglycerides, fasting glucose or fasting insulin, insulin resistance (HOMA-IR) or  
406 insulin sensitivity (QUICKI) after 4 weeks of isocaloric TRE or normal diet in elite cyclists.

407

408 In regards to muscular strength and cardio respiratory fitness, seven studies have  
409 examined the effect of time restricted eating combined with exercise.<sup>29,31,34-36</sup> Correia et al.<sup>29</sup>

410 reported an increase in explosive upper body strength from baseline in TRE and normal diet  
411 groups. A significant time by diet interaction for peak force and peak dynamic bench press  
412 throw favoring the TRE group. The normal diet group also increased their explosive upper body  
413 strength over time, produced greater improvements over time compared to TRE (time by diet  
414 interaction) for squat jump peak force, countermovement jump peak force, countermovement  
415 jump height and isometric bench press. Moro et al.<sup>31</sup> reported significant increases in leg press  
416 and hip sled in both the TRE combined with resistance training and the normal diet combined  
417 with resistance training groups. Bench press and leg press increased over time in both the TRE  
418 and normal diet groups with no difference between groups.<sup>42</sup> Tinsley et al.<sup>34</sup> also reported an  
419 increase in hip sled, hip sled endurance and bench press when combining resistance training with  
420 both TRE and normal diet after 8 weeks in trained healthy males. Brady et al.<sup>36</sup> reported no  
421 significant changes in fixed blood lactate concentration, heart rate at fixed blood lactate  
422 concentration, and %HR max or  $VO_{2max}$  in either group after 8 weeks. Tinsley et al.<sup>35</sup> reported an  
423 increase in maximum strength and muscular performance (countermovement vertical jump,  
424 mechanized squat, and 1 repetition max and repetitions to failure of bench press and hip sled)  
425 when combining resistance training with TRE or normal 12-h diet, independent of HMB  
426 supplementation in resistance trained females. Martinez-Rodriguez et al.<sup>39</sup> reported an increase  
427 from baseline countermovement vertical jump height in the TRE combined with HIIT group.  
428 Additionally, there was a group interaction at week 16 between the combination group and HIIT  
429 alone group.

430

431 Summary of findings: In individuals with overweight or obesity TRE combined with  
432 exercise produced significant decreases in body weight of 2-4% after 7-16 weeks. Body fat mass

433 and waist circumference also seem to decrease significantly. Changes in fat free mass changes  
434 were inconsistent and may depend on intensity or volume of aerobic or resistance training. In  
435 normal weight, trained individuals body weight appears unchanged in trials prescribed a calorie  
436 goal for weight maintenance and decreases in trials with an ad libitum eating window. However,  
437 fat mass does appear to decrease significantly when combining TRE and exercise in lean  
438 individuals. LDL cholesterol and blood pressure remained largely unchanged, however TRE  
439 combined with exercise may increase HDL cholesterol and decrease triglycerides in normal  
440 weight individuals. Glucoregulatory factors also appear to be unaffected by TRE and exercise  
441 independent of BMI category. However, one long term follow-up did report significantly  
442 improved glucose, insulin, and insulin resistance in lean trained males,<sup>42</sup> indicating these changes  
443 may improve over longer time periods. It appears that if caloric intake is adequate, improvements  
444 in explosive strength, muscular endurance,  $V_{O_{2peak}}$  can still be achieved with TRE. It is unclear  
445 how TRE may influence adaptations to aerobic or strength training in untrained individuals or  
446 those with overweight or obesity.

## 447 **DISCUSSION**

448 Over 75% of Americans have either overweight or obesity.<sup>1</sup> This is startling as obesity is  
449 associated co-morbidities such as heart disease, cancer, stroke, and diabetes.<sup>1</sup> One in five adults  
450 in the United States is inactive, which paired with rates of obesity, greatly increase risk of lower  
451 quality of life, mental health issues, comorbidities and mortality.<sup>6</sup> While combining different  
452 forms of intermittent fasting with exercise has shown favorable effects on body weight and body  
453 fat, improvements in cardiovascular and metabolic risk were not consistent. The results are  
454 limited by 1) sample size, 2) intervention length, 3) inclusion criteria, 4) objective measures of  
455 diet and exercise adherence, and 5) diversity of sample population.

456

457 First, small sample size is a considerable limitation of many of the studies that were  
458 reviewed here. Sixteen of the 23 trials reviewed were pilot studies with less than 50 participants,  
459 which were then randomized into 2-4 groups. This indicates that most of the trials presented here  
460 may be underpowered to report primary and secondary outcomes. Second, current trials lack  
461 long-term testing and follow-up. Of the 23 trials reported in this review, 18 were short term (4-12  
462 weeks), 2 were mid-term (16 weeks), and 3 were long term (52 weeks). Two long-term (52  
463 week) studies combining 5:2 with exercise did report significant improvements in blood  
464 pressure, HDL cholesterol, and triglycerides. At a follow-up at one year after an 8-week study of  
465 TRE combined with resistance training, Moro et al.<sup>42</sup> reported significant improvements in  
466 cardiometabolic markers including a significant time by diet interaction in insulin resistance.  
467 This may suggest that prior interventions were not long enough to achieve optimal effects. Third,  
468 participants included in the current breadth of work, even those with obesity, were metabolically  
469 healthy at baseline (being excluded if they had hypertension, dyslipidemia or pre-diabetes). Thus,  
470 while participants may have benefited from decreases in body weight and body fat mass, the  
471 potential impact on cardiometabolic risk may have been masked by a floor effect related to the  
472 inclusion criteria. Specifically, of the 13 trials combining TRE with exercise, only three<sup>33,37,38</sup>  
473 examined individuals with overweight or obesity and two<sup>33,37</sup> included people who were not  
474 already physically active. Trained individuals are more likely to be euglycemic to begin with,  
475 given the direct influence of exercise on glucoregulation.<sup>44-46</sup> Thus, future studies should focus  
476 on recruitment of higher risk groups, such as those with prediabetes and untrained individuals.  
477 Fourth, objective data on adherence and compliance to both intermittent fasting and exercise  
478 interventions are lacking. Currently, food diaries and other self-report techniques are utilized to



479 monitor adherence and compliance to different forms of intermittent fasting. It is well established  
480 that individuals under-report energy intake and selectively report foods that are considered to be  
481 “healthy” or socially acceptable.<sup>47</sup> It will be important for future studies to explore more  
482 objective measures of adherence to these fasting diets such as continuous glucose monitors. As  
483 for the exercise interventions, only eleven<sup>16,20,21,23,24,31,33-35,37,39</sup> of the reviewed trials either  
484 supervised all exercise or utilized wearables (Actigraph, Pensacola FL or Garmin, Olathe KS) to  
485 monitor adherence and compliance. To determine efficacy of these behavioral interventions,  
486 high-quality adherence and compliance data are essential. Lastly, racial, ethnic, sex, and age  
487 diversity should be considered in future work to increase the external validity of the outcomes.  
488 Current data in 5:2 or ADF combined with exercise do include both males and females as well as  
489 those aged 18-65 years or older, however, this is a stark contrast when examining the current data  
490 in TRE combined with exercise. Of the 13 trials presented combining TRE with exercise, only  
491 five<sup>33,35,37-39,48</sup> included women and two<sup>30,37</sup> included those aged 45 years or older. The remaining  
492 trials are focused in lean, young, male athletes which is not representative of the U.S. population.  
493 While the trials described here span the globe including the US, Brazil, Australia, New Zealand,  
494 China, Korea, Germany, Italy, Portugal, Norway, Ireland and Spain, most of these trials have  
495 been in European predominantly White countries. Of the seven<sup>23,24,28,30,34,35,37</sup> trials in the US  
496 only two<sup>23,24</sup> reported on race and ethnicity. Due to the impact of social determinants of health,  
497 including race and ethnicity, on obesity and cardiometabolic risk it is imperative that research  
498 includes those from underrepresented backgrounds.

499         Due to the above limitations, future randomized controlled trials should deliver longer  
500 interventions ( $\geq 24$  weeks) with larger, diverse sample sizes to assess efficacy of intermittent  
501 fasting combined with exercise. Individuals who are young, healthy, active, and lean do not have

502 the same heightened risk and are thus unlikely to improve cardiometabolic health based on floor  
503 effects. The benefits of intermittent fasting may be greatest for those with obesity, overweight,  
504 and/or cardiometabolic risk, and thus more research is needed in these groups. Future studies  
505 should utilize tools to measure adherence and compliance to both the diet and exercise programs  
506 and explore ways to obtain objective data. Lastly, it is imperative that researchers include both  
507 males and females and persons of diverse racial and ethnic backgrounds and across the lifespan  
508 in interventions combining diet and exercise, consistent with recent policy advancements and  
509 position statements from the U.S. National Institutes of Health and related organizations.

## 510 **CONCLUSION**

511 The 5:2 diet, ADF, and TRE offer accessible and sustainable alternatives to traditional CR. When  
512 combined with different modalities of exercise, these diets can reduce body weight and fat mass.  
513 Although there is uncertain impact on chronic disease risk, there is some evidence to suggest that  
514 benefits may emerge in higher-powered and longer interventions. Training adaptations are still  
515 possible when combining any form of intermittent fasting with exercise. Ongoing research is  
516 needed to test the benefits of combined interventions in diverse populations.

517 **REFERENCES**

- 518 1. Prevention CfDca. Adult Obesity Facts. US Department of Health and Human Services.  
519 Accessed 03/05/2018, 2018. <https://www.cdc.gov/obesity/data/adult.html>
- 520 2. Trepanowski JF, Kroeger CM, Barnosky A, et al. Effects of alternate-day fasting or daily  
521 calorie restriction on body composition, fat distribution, and circulating adipokines: Secondary  
522 analysis of a randomized controlled trial. *Clin Nutr*. Dec 2017;doi:10.1016/j.clnu.2017.11.018
- 523 3. Harvie MN, Pegington M, Mattson MP, et al. The effects of intermittent or continuous  
524 energy restriction on weight loss and metabolic disease risk markers: a randomized trial in  
525 young overweight women. *Int J Obes (Lond)*. May 2011;35(5):714-27. doi:10.1038/ijo.2010.171
- 526 4. Varady KA, Cienfuegos S, Ezpeleta M, Gabel K. Cardiometabolic Benefits of Intermittent  
527 Fasting. *Annu Rev Nutr*. Oct 11 2021;41:333-361. doi:10.1146/annurev-nutr-052020-041327
- 528 5. Ezpeleta M, Cienfuegos S, Lin S, et al. Time-restricted eating: Watching the clock to treat  
529 obesity. *Cell Metabolism*. 2024/01/03/ 2024;doi:<https://doi.org/10.1016/j.cmet.2023.12.004>
- 530 6. Powell KE, King AC, Buchner DM, et al. The Scientific Foundation for the Physical Activity  
531 Guidelines for Americans, 2nd Edition. *J Phys Act Health*. Dec 17 2018:1-11.  
532 doi:10.1123/jpah.2018-0618
- 533 7. Oh H, Saquib N, Ochs-Balcom HM, et al. Recreational Physical Activity, Sitting, and  
534 Androgen Metabolism among Postmenopausal Women in the Women's Health Initiative  
535 Observational Study. *Cancer Epidemiol Biomarkers Prev*. Jan 2022;31(1):97-107.  
536 doi:10.1158/1055-9965.Epi-21-0809

- 537 8. Willoughby D, Hewlings S, Kalman D. Body Composition Changes in Weight Loss:  
538 Strategies and Supplementation for Maintaining Lean Body Mass, a Brief Review. *Nutrients*. Dec  
539 3 2018;10(12)doi:10.3390/nu10121876
- 540 9. Hansen D, Dendale P, Berger J, van Loon LJ, Meeusen R. The effects of exercise training  
541 on fat-mass loss in obese patients during energy intake restriction. *Sports Med*. 2007;37(1):31-  
542 46. doi:10.2165/00007256-200737010-00003
- 543 10. Chow ZS, Moreland AT, Macpherson H, Teo WP. The Central Mechanisms of Resistance  
544 Training and Its Effects on Cognitive Function. *Sports Med*. Dec 2021;51(12):2483-2506.  
545 doi:10.1007/s40279-021-01535-5
- 546 11. Bellicha A, van Baak MA, Battista F, et al. Effect of exercise training on weight loss, body  
547 composition changes, and weight maintenance in adults with overweight or obesity: An  
548 overview of 12 systematic reviews and 149 studies. *Obes Rev*. Jul 2021;22 Suppl 4(Suppl  
549 4):e13256. doi:10.1111/obr.13256
- 550 12. Ryan DH, Yockey SR. Weight Loss and Improvement in Comorbidity: Differences at 5%,  
551 10%, 15%, and Over. *Curr Obes Rep*. Jun 2017;6(2):187-194. doi:10.1007/s13679-017-0262-y
- 552 13. Das JK, Banskota N, Candia J, et al. Calorie restriction modulates the transcription of  
553 genes related to stress response and longevity in human muscle: The CALERIE study. *Aging cell*.  
554 Oct 12 2023:e13963. doi:10.1111/accel.13963
- 555 14. Das SK, Gilhooly CH, Golden JK, et al. Long-term effects of 2 energy-restricted diets  
556 differing in glycemic load on dietary adherence, body composition, and metabolism in CALERIE:  
557 a 1-y randomized controlled trial. *Am J Clin Nutr*. Apr 2007;85(4):1023-30.  
558 doi:10.1093/ajcn/85.4.1023

- 559 15. Keenan S, Cooke MB, Chen WS, Wu S, Belski R. The Effects of Intermittent Fasting and  
560 Continuous Energy Restriction with Exercise on Cardiometabolic Biomarkers, Dietary  
561 Compliance, and Perceived Hunger and Mood: Secondary Outcomes of a Randomised,  
562 Controlled Trial. *Nutrients*. 2022;14(15)doi:10.3390/nu14153071
- 563 16. Batitucci G, Faria Junior EV, Nogueira JE, et al. Impact of Intermittent Fasting Combined  
564 With High-Intensity Interval Training on Body Composition, Metabolic Biomarkers, and Physical  
565 Fitness in Women With Obesity. *Front Nutr*. 2022;9:884305. doi:10.3389/fnut.2022.884305
- 566 17. Kang J, Shi X, Fu J, Li H, Ma E, Chen W. Effects of an Intermittent Fasting 5:2 Plus Program  
567 on Body Weight in Chinese Adults with Overweight or Obesity: A Pilot Study. *Nutrients*.  
568 2022;14(22)doi:10.3390/nu14224734
- 569 18. Hottenrott K, Werner T, Hottenrott L, Meyer TP, Vormann J. Exercise Training,  
570 Intermittent Fasting and Alkaline Supplementation as an Effective Strategy for Body Weight Loss:  
571 A 12-Week Placebo-Controlled Double-Blind Intervention with Overweight Subjects. *Life (Basel)*.  
572 2020;10(5)doi:10.3390/life10050074
- 573 19. Headland ML, Clifton PM, Keogh JB. Effect of intermittent compared to continuous  
574 energy restriction on weight loss and weight maintenance after 12 months in healthy  
575 overweight or obese adults. *Int J Obes (Lond)*. Oct 2019;43(10):2028-2036. doi:10.1038/s41366-  
576 018-0247-2
- 577 20. Jospe MR, Roy M, Brown RC, et al. Intermittent fasting, Paleolithic, or Mediterranean  
578 diets in the real world: exploratory secondary analyses of a weight-loss trial that included choice  
579 of diet and exercise. *Am J Clin Nutr*. Mar 1 2020;111(3):503-514. doi:10.1093/ajcn/nqz330

- 580 21. Cooke MB, Deasy W, Ritenis EJ, Wilson RA, Stathis CG. Effects of Intermittent Energy  
581 Restriction Alone and in Combination with Sprint Interval Training on Body Composition and  
582 Cardiometabolic Biomarkers in Individuals with Overweight and Obesity. *International journal of*  
583 *environmental research and public health*. Jun 29 2022;19(13)doi:10.3390/ijerph19137969
- 584 22. Keenan SJ, Cooke MB, Hassan EB, et al. Intermittent fasting and continuous energy  
585 restriction result in similar changes in body composition and muscle strength when combined  
586 with a 12 week resistance training program. *Eur J Nutr*. 2022;61(4):2183-2199.  
587 doi:10.1007/s00394-022-02804-3
- 588 23. Bhutani S, Klempel MC, Kroeger CM, Trepanowski JF, Varady KA. Alternate day fasting  
589 and endurance exercise combine to reduce body weight and favorably alter plasma lipids in  
590 obese humans. *Obesity (Silver Spring)*. Jul 2013;21(7):1370-9. doi:10.1002/oby.20353
- 591 24. Ezpeleta M, Gabel K, Cienfuegos S, et al. Effect of alternate day fasting combined with  
592 aerobic exercise on non-alcoholic fatty liver disease: A randomized controlled trial. *Cell Metab*.  
593 Jan 3 2023;35(1):56-70 e3. doi:10.1016/j.cmet.2022.12.001
- 594 25. Cho AR, Moon JY, Kim S, et al. Effects of alternate day fasting and exercise on cholesterol  
595 metabolism in overweight or obese adults: A pilot randomized controlled trial. *Metabolism*. Apr  
596 2019;93:52-60. doi:10.1016/j.metabol.2019.01.002
- 597 26. Oh M, Kim S, An KY, et al. Effects of alternate day calorie restriction and exercise on  
598 cardio-metabolic risk factors in overweight and obese adults: an exploratory randomized  
599 controlled study. *BMC public health*. Sep 15 2018;18(1):1124. doi:10.1186/s12889-018-6009-1

- 600 27. Moro T, Tinsley G, Longo G, et al. Time-restricted eating effects on performance, immune  
601 function, and body composition in elite cyclists: a randomized controlled trial. *J Int Soc Sports*  
602 *Nutr.* Dec 11 2020;17(1):65. doi:10.1186/s12970-020-00396-z
- 603 28. Richardson CE, Tovar AP, Davis BA, Van Loan MD, Keim NL, Casazza GA. An Intervention  
604 of Four Weeks of Time-Restricted Eating (16/8) in Male Long-Distance Runners Does Not Affect  
605 Cardiometabolic Risk Factors. *Nutrients.* Feb 16 2023;15(4)doi:10.3390/nu15040985
- 606 29. Correia JM, Santos PDG, Pezarat-Correia P, Minderico CS, Infante J, Mendonca GV. Effect  
607 of Time-Restricted Eating and Resistance Training on High-Speed Strength and Body  
608 Composition. *Nutrients.* Jan 6 2023;15(2)doi:10.3390/nu15020285
- 609 30. Waldman HS, Witt CR, Grozier CD, McAllister MJ. A self-selected 16:8 time-restricted  
610 eating quasi-experimental intervention improves various markers of cardiovascular health in  
611 middle-age male cyclists. *Nutrition.* May 23 2023;113:112086. doi:10.1016/j.nut.2023.112086
- 612 31. Moro T, Tinsley G, Bianco A, et al. Effects of eight weeks of time-restricted feeding (16/8)  
613 on basal metabolism, maximal strength, body composition, inflammation, and cardiovascular  
614 risk factors in resistance-trained males. *J Transl Med.* Oct 2016;14(1):290. doi:10.1186/s12967-  
615 016-1044-0
- 616 32. McAllister MJ, Pigg BL, Renteria LI, Waldman HS. Time-restricted feeding improves  
617 markers of cardiometabolic health in physically active college-age men: a 4-week randomized  
618 pre-post pilot study. *Nutr Res.* Mar 2020;75:32-43. doi:10.1016/j.nutres.2019.12.001
- 619 33. Haganes KL, Silva CP, Eyjólfssdóttir SK, et al. Time-restricted eating and exercise training  
620 improve HbA1c and body composition in women with overweight/obesity: A randomized  
621 controlled trial. *Cell Metab.* Oct 4 2022;34(10):1457-1471.e4. doi:10.1016/j.cmet.2022.09.003

- 622 34. Tinsley GM, Forsse JS, Butler NK, et al. Time-restricted feeding in young men performing  
623 resistance training: A randomized controlled trial. *Eur J Sport Sci.* Mar 2017;17(2):200-207.  
624 doi:10.1080/17461391.2016.1223173
- 625 35. Tinsley GM, Moore ML, Graybeal AJ, et al. Time-restricted feeding plus resistance  
626 training in active females: a randomized trial. *Am J Clin Nutr.* Sep 1 2019;110(3):628-640.  
627 doi:10.1093/ajcn/nqz126
- 628 36. Brady AJ, Langton HM, Mulligan M, Egan B. Effects of 8 wk of 16:8 Time-restricted Eating  
629 in Male Middle- and Long-Distance Runners. *Med Sci Sports Exerc.* Mar 1 2021;53(3):633-642.  
630 doi:10.1249/MSS.0000000000002488
- 631 37. Kotarsky CJ, Johnson NR, Mahoney SJ, et al. Time-restricted eating and concurrent  
632 exercise training reduces fat mass and increases lean mass in overweight and obese adults.  
633 *Physiological reports.* May 2021;9(10):e14868. doi:10.14814/phy2.14868
- 634 38. Isenmann E, Dissemond J, Geisler S. The Effects of a Macronutrient-Based Diet and Time-  
635 Restricted Feeding (16:8) on Body Composition in Physically Active Individuals-A 14-Week  
636 Randomised Controlled Trial. *Nutrients.* Sep 6 2021;13(9)doi:10.3390/nu13093122
- 637 39. Martínez-Rodríguez A, Rubio-Arias JA, García-De Frutos JM, Vicente-Martínez M,  
638 Gunnarsson TP. Effect of High-Intensity Interval Training and Intermittent Fasting on Body  
639 Composition and Physical Performance in Active Women. *International journal of environmental*  
640 *research and public health.* Jun 14 2021;18(12)doi:10.3390/ijerph18126431
- 641 40. Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care.*  
642 Jun 2004;27(6):1487-95.



- 643 41. Chen H, Sullivan G, Quon MJ. Assessing the Predictive Accuracy of QUICKI as a Surrogate  
644 Index for Insulin Sensitivity Using a Calibration Model. *Diabetes*. 2005;54(7):1914-1925.  
645 doi:10.2337/diabetes.54.7.1914
- 646 42. Moro T, Tinsley G, Pacelli FQ, Marcolin G, Bianco A, Paoli A. Twelve Months of Time-  
647 restricted Eating and Resistance Training Improves Inflammatory Markers and Cardiometabolic  
648 Risk Factors. *Med Sci Sports Exerc*. Dec 1 2021;53(12):2577-2585.  
649 doi:10.1249/mss.0000000000002738
- 650 43. Witt CR, Grozier CD, Killen LG, Renfroe LG, O'Neal EK, Waldman HS. A Self-Selected 16:8  
651 Time-Restricted Eating Protocol Improves Fat Oxidation Rates, Markers of Cardiometabolic  
652 Health, and 10-km Cycling Performance in Middle-Age Male Cyclists. *J Strength Cond Res*. May 1  
653 2023;37(5):1117-1123. doi:10.1519/jsc.0000000000004353
- 654 44. Ploug T, van Deurs B, Ai H, Cushman SW, Ralston E. Analysis of GLUT4 distribution in  
655 whole skeletal muscle fibers: identification of distinct storage compartments that are recruited  
656 by insulin and muscle contractions. *J Cell Biol*. Sep 21 1998;142(6):1429-46.  
657 doi:10.1083/jcb.142.6.1429
- 658 45. Kranjic GN, Cameron-Smith D, Hargreaves M. Acute exercise and GLUT4 expression in  
659 human skeletal muscle: influence of exercise intensity. *J Appl Physiol (1985)*. Sep  
660 2006;101(3):934-7. doi:10.1152/jappphysiol.01489.2005
- 661 46. Mikines KJ, Sonne B, Farrell PA, Tronier B, Galbo H. Effect of physical exercise on  
662 sensitivity and responsiveness to insulin in humans. *Am J Physiol*. Mar 1988;254(3 Pt 1):E248-  
663 59. doi:10.1152/ajpendo.1988.254.3.E248

- 664 47. Ortega RM, Perez-Rodrigo C, Lopez-Sobaler AM. Dietary assessment methods: dietary  
665 records. *Nutr Hosp*. Feb 26 2015;31 Suppl 3:38-45. doi:10.3305/nh.2015.31.sup3.8749
- 666 48. Martinez-Gonzalez MA, Fernandez-Jarne E, Serrano-Martinez M, Wright M, Gomez-  
667 Gracia E. Development of a short dietary intake questionnaire for the quantitative estimation of  
668 adherence to a cardioprotective Mediterranean diet. *Eur J Clin Nutr*. Nov 2004;58(11):1550-2.  
669 doi:10.1038/sj.ejcn.1602004
- 670

**Table 1. Designs: Design, participant characteristics, and intervention descriptions of human trials of intermittent fasting combined with exercise.**

Reference	Sample Size	Participant characteristics	Diet length	Design	Intervention arms	Exercise intervention specifics
<b>5:2 Diet</b>						
Batitucci 2022 <sup>16</sup>	n=36	Female Obesity	8 weeks	<b>RT<sup>a</sup>: Parallel- arm</b>	1. 5:2: Fast day (600kcal) Feast day (ad libitum) 2. 5:2 + Exercise 3. Exercise	HIIT <sup>b</sup> 3 days/week
Keenan 2021, 2022 <sup>15,22</sup>	n=44 n=34 analyzed	Male/Female normal weight, overweight, obese recreationally active	12 weeks	<b>RT<sup>a</sup>: Parallel- arm</b>	1. 5:2: Fast day (-30% TEE <sup>c</sup> ) + resistance training. Fasting meals provided. 2. CR <sup>d</sup> (-20%+) + resistance training	2 supervised resistance training sessions + 1 unsupervised aerobic/resistance training session
Kang 2022 <sup>17</sup>	n= 131	Male/Female Overweight or Obesity	12 weeks	<b>RT<sup>a</sup>: Parallel- arm</b>	1. 5:2: Fast day (30% TEE <sup>c</sup> ) Feast day (70% TEE <sup>c</sup> ) 2. CR <sup>d</sup> (70% TEE <sup>c</sup> ) 3. High protein meal replacement (70% TEE <sup>c</sup> provided)	150-300m physical activity
Hottenrott 2020 <sup>18</sup>	n=80 n=68 analyzed	Male/Female Overweight "Healthy"	12 weeks	<b>RCT<sup>e</sup>: Parallel- arm</b>	1. 5:2: Fast day (F: 400, M:600 kcal) Feast day (ad libitum) Alkaline Supplement 2. 5:2: Fast day (F: 500, M:600 kcal) Feast day (ad libitum) Placebo 3. Ad libitum Alkaline Supplement 4. Ad libitum Placebo	Exercise in all groups: 30-60 minutes of running and 20 minutes of resistance training 3-4 days/week
Cooke 2022 <sup>21</sup>	n=34	Male/Female Overweight or obesity	16 weeks	<b>RT<sup>a</sup>: Parallel- arm Per protocol</b>	1. 5:2 (ad libitum) 2. Sprint interval training 3. 5:2 + sprint interval training	Sprint interval training 3 days/week 4 × 20 s work followed by 40 s of active rest
Headland 2019 <sup>19</sup>	n=332 n=124 analyzed	Male/Female Overweight or obesity	52 weeks	<b>RT<sup>a</sup>: Parallel- arm</b>	1. 5:2: Fast day (F: 500, M:600 kcal) Feast day (ad libitum) 2. Week-on, week-off (F: 1000, M: 120 0kcal/d) Week-off (ad libitum) 3. CR <sup>d</sup> (-30% TEE <sup>c</sup> )	All groups: Increase to 10,000 steps
Jospe 2020 <sup>20</sup>	n=250 n=171 analyzed	Male/Female, Overweight or obesity	52 weeks	<b>Parallel- arm non-randomized Per protocol</b>	1. Mediterranean Diet 2. 5:2 3. Paleolithic Self-selected diet arm	Choice of standard physical activity recommendations OR home based HIIT <sup>b</sup>
<b>Alternate day fasting</b>						

Cho 2019 <sup>25,26</sup>	n=100 n=31/33 analyzed	Male/Female Overweight or obesity	8 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. ADF <sup>f</sup> +Exercise 2. ADF <sup>f</sup> 3. Ex 4. Control	Resistance and aerobic training 3 days/week. First week only supervised.
Bhutani 2013 <sup>23</sup>	N=64	Male/Female Obesity	12 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. ADF <sup>f</sup> +Exercise 2. ADF <sup>f</sup> 3. Exercise 4. Control	Aerobic activity 3 days/week, supervised
Ezpeleta 2023 <sup>24</sup>	n=80	Male/Female Obesity non-alcoholic fatty liver disease	12 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. ADF <sup>f</sup> 2. Exercise alone 3. ADF <sup>f</sup> +Exercise 4. Control	Aerobic activity 5 days/week, supervised
<b>Time restricted eating</b>						
Moro 2020 <sup>27</sup>	n=16	Male elite cyclists	4 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> (8h, 10am-6PM) 2. Normal Diet (12h) Isocaloric (7d diet plan) 3 meals + one snack	500 km/week over 6 sessions
Richardson 2023 <sup>28</sup>	n=24 n=15 analyzed	Male endurance trained runners	4 weeks, 2-week washout	<b>RCT<sup>o</sup>: Crossover</b>	1. TRE <sup>g</sup> (8h, self-selected) 2. Normal Diet (12h) Isocaloric	Maintain current training
Correia 2023 <sup>29</sup>	n= 18	Male healthy trained	30 days	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> + resistance training 2. Normal diet (12h) + resistance training	Resistance training 3 days/week, 1 time/week supervised
Waldman 2023 <sup>30,43</sup>	n=15 n=12 analyzed	Male cyclists	4 weeks	<b>Single arm</b>	1. TRE <sup>g</sup> (16h self-selected)	150 minutes per week
Haganes 2022 <sup>33</sup>	n=131	Male/Female Overweight or Obesity	7 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> 2. HIIT <sup>g</sup> 3. TRE <sup>g</sup> and HIIT <sup>b</sup> 4. Control	HIIT <sup>b</sup> (running) 3 days per week, supervised and wearable utilized
Moro 2016 <sup>31</sup>	n=34	Male healthy trained	8 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> (4h 4d/wk) + resistance training 2. Normal diet (12h) + resistance training Weight maintenance calorie goal	Resistance training 3 non-consecutive days/week
Moro 2021 <sup>42</sup>	n=20	Male healthy	Follow-up at 52 weeks <sup>31</sup>	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> (1-9PM) 2. Normal diet (12h)	Resistance training 3 non-consecutive days/week, supervised
Tinsley 2017 <sup>34</sup>	n=18	Male healthy trained	8 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> (1-8PM) + resistance training 2. Normal diet (12h) + resistance training Weight maintenance calorie goal	Resistance training 3 days/week on non-fasting days
Brady 2021 <sup>36</sup>	n=23 n=17 analyzed	Male middle/ long distance runners	8 weeks	<b>RCT<sup>o</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> (8h) 2. Control	Maintain habitual running, wearable utilized
Tinsley 2019 <sup>35</sup>	n=40	Female Resistance trained	8 weeks	<b>RCT<sup>o</sup>: Placebo controlled</b>	1. TRE <sup>g</sup> (8h) + resistance training 2. TRE <sup>g</sup> (8h) + supplement + resistance training 3. Normal diet (12h) + resistance training	Resistance training 3 non-consecutive days/week, supervised and wearable utilized

Kotarsky 2021 <sup>37</sup>	n=28 n=21 analyzed	Male/Female Overweight or obesity	8 weeks	<b>RCT<sup>e</sup>: Parallel- arm</b>	1. TRE <sup>g</sup> (12- 8PM, ad libitum) 2. Control (normal eating)	300 minutes of moderate or 150 vigorous aerobic and resistance training per week, resistance training supervised, wearable utilized
Iseemann 2021 <sup>38</sup>	n=35	Male/Female Overweight or obesity	2 weeks familiarization n 8 weeks intervention	<b>RT<sup>a</sup> : Parallel- arm Per protocol</b>	1. TRE <sup>g</sup> (12-8PM, ad libitum but macronutrient goal breakdown given) 2. MBD <sup>h</sup> (80% unprocessed, 20% could be processed, chose foods based on the Nutri-score scale)	Two group training sessions per week, unsupervised, gym attendance checked
Martínez-Rodríguez 2021 <sup>39</sup>	n=14	Female active normal weight	16 weeks	<b>RT<sup>a</sup> : Crossover</b>	1. TRE <sup>g</sup> + HIIT <sup>b</sup> 2. HIIT <sup>b</sup>	HIIT <sup>b</sup> 3 days /week (40 minutes) 3 × 10 repetitions of 30 s of aerobic exercises all out alternated with 30 s of passive recovery, supervised

<sup>a</sup>RT: Randomized trial, <sup>b</sup>HIIT: high intensity interval training, <sup>c</sup>TEE: total energy expenditure, <sup>d</sup>CR: Calorie restriction, <sup>e</sup>RCT: Randomized controlled trial, <sup>f</sup>ADF: Alternate day fasting, <sup>g</sup>TRE: time restricted eating, <sup>h</sup>MBD: macronutrient-based diet

**Table 2. Results: Effects of intermittent fasting combined with exercise on body weight, body composition, cardiometabolic markers and muscular strength and cardiorespiratory fitness.**

Reference	Participants	Diet length	Intervention Groups	Body weight	Body composition			Blood pressure	Plasma lipids			Glucoregulatory factors			Performance/ Strength
					FM <sup>a</sup>	FFM <sup>b</sup>	VF <sup>c</sup>		LDL <sup>d</sup>	HDL <sup>e</sup>	TG <sup>f</sup>	Fasting Glucose	Fasting Insulin	IR <sup>g</sup> /IS <sup>h</sup> /A1c <sup>i</sup>	
5:2 Diet															
Batitucci 2022 <sup>16</sup>	n=36	8-week	1. 5:2 2. 5:2 + Ex <sup>j</sup> 3. Ex <sup>j</sup>	1. ∅ 2. ↓2% 3. ∅	1. ∅ 2. ↓ 3. ↓	1. ∅ 2. ↑ 3. ↑	1. ∅ 2. 3. ↓	--	--	--	--	--	--	--	1. ∅ 2. ↑ walking, strength, HRmax <sup>k</sup> , V02peak <sup>l</sup> 3. ↑ walking, strength, HRmax <sup>k</sup> , V02peak <sup>l</sup>
Keenan 2021, 2022 <sup>15,22</sup>	n=44 n=34 analyzed	12-week	1. 5:2 + Ex <sup>j</sup> 2. CR <sup>m</sup> + Ex <sup>j</sup>	1. ↓4% 2. ↓5%	1. ↓ 2. ↓	1. ↑ 2. ↑	--	--	1. ↓† 2. ↓	1. ↓ 2. ↓	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅ -	1. ↑ upper and lower body strength 2. ↑ muscle surface area, upper and lower body strength
Kang 2022 <sup>17</sup>	n= 131	12-week	1. 5:2 2. CR <sup>m</sup> 3. High protein meal replacement	1. ↓9%† 2. ↓5% 3. ↓9%†	1. ↓ 2. ↓ 3. ↓	1. ↑ 2. ↑ 3. ↑	--	--	--	--	--	--	--	--	--
Hottenrott 2020 <sup>18</sup>	n=80 n=68 analyzed	12-week	1. 5:2 + Alkaline supplement 2. 5:2 Placebo 3. Ad Lib <sup>n</sup> Alkaline Supplement 4. Ad lib <sup>n</sup> Placebo	1. ↓8kg† 2. ↓6kg 3. ↓6kg 4. ↓3kg	1. ↓† 2. ↓ 3. ↓ 4. ↓	--	1. ↓† 2. ↓ 3. ↓ 4. ↓	--	--	--	--	--	--	--	1. ↑† running velocity 2. ∅ 3. ∅ 4. ∅
Cooke 2022 <sup>21</sup>	n=34	16-week	1. 5:2 2. SIT <sup>o</sup> 3. 5:2 + SIT <sup>o</sup>	1. ↓† 2. ∅ 3. ↓†	1. ↓† 2. ∅ 3. ↓†	1. ↓ 2. ↓ 3. ↓	1. ↓ 2. ↓ 3. ↓ WC <sup>p</sup>	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	1. ∅ IR <sup>g</sup> 2. ∅ IR <sup>g</sup> 3. ∅ IR <sup>g</sup>	1. ∅ 2. ↑† V02peak <sup>l</sup> 3. ↑† V02peak <sup>l</sup>	
Headland 2019 <sup>19</sup>	n=332 n=124 analyzed	52 weeks	1. 5:2 2. Week-on, week-off 3. CR <sup>m</sup>	1. ↓6% 2. ↓6% 3. ↓8%	1. ↓ 2. ↓ 3. ↓	1. ↓ 2. ↓ 3. ↓	--	--	1. ∅ 2. ∅ 3. ∅	1. ↑ 2. ↑ 3. ↑	1. ↓ 2. ↓ 3. ↓	1. ∅ 2. ∅ 3. ∅	--	--	--
Jospe 2020 <sup>20</sup>	n=250	52 weeks	1. Mediterranean	1. ↓ 3% 2. ↓4%	1. ↓ 2. ↓	--	1. ↓ 2. ↓	1. ↓SBP <sup>q</sup> ↓DBP <sup>r</sup>	1. ↓ 2. ∅	1. ∅ 2. ↑	1. ∅ 2. ∅	--	--	1. ↓A1c <sup>l</sup> 2. ∅	--

	n=171 analyzed		Diet 2. 5:2 3. Paleolithic	3. ∅	3. ∅		3. ∅	2. ↓ SBP <sup>q</sup> ↓ DBP <sup>r</sup> 3. ∅ SBP <sup>q</sup> ↓ DBP <sup>r</sup>	3. ↓	3. ∅	3. ∅			3. ∅	
Alternate Day Fasting															
Cho 2019 <sup>25,26</sup>	n=100 n=31/33 analyzed	8 weeks	1. ADF <sup>s</sup> + Ex <sup>j</sup> 2. ADF <sup>s</sup> 3. Ex <sup>j</sup> 4. Control	1. ↓4%† 2. ↓3%† 3. ∅ 4. ∅	1. ↓† 2. ↓† 3. ↓ 4. ↓	1. ↓ 2. ∅ 3. ∅ 4. ∅	1. ↓ 2. ↓ 3. ↓ 4. ∅	--	1. ∅ 2. ∅ 3. ↑ 4. ∅	1. ↑ 2. ∅ 3. ↑ 4. ↑	1. ↓ 2. ↑† 3. ↓† 4. ↑†	1. ↓ 2. ↓ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ↑	1. ∅ IR <sup>o</sup> 2. ∅ IR <sup>o</sup> 3. ∅ IR <sup>o</sup> 4. ↑ IR <sup>o</sup>	1. ↑ V02peak <sup>l</sup> , ↑ Mets min/wk, muscle strength (chest press, shoulder press, lat pull) 2. ↓ Chest press 3. ↑ Muscle strength (chest press, shoulder press, lat pull) 4. ∅
Bhutani 2013 <sup>23</sup>	N=64	12 weeks	1. ADF <sup>s</sup> + Ex <sup>j</sup> 2. ADF <sup>s</sup> 3. Ex <sup>j</sup> 4. Control	1. ↓7%† 2. ↓3% 3. 1% ↓ 4. ∅	1. ↓† 2. ↓ 3. ∅ 4. ∅	1. ∅ 2. ↓ 3. ∅ 4. ∅	1. ↓† 2. ↓ 3. ↓ 4. ∅	1. ∅ 2. ↓ SBP <sup>q</sup> ↓ DBP <sup>r</sup> 3. ∅ 4. ∅	1. ↓ 2. ∅ 3. ∅ 4. ∅	1. ↑† 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ IR <sup>o</sup> 2. ∅ IR <sup>o</sup> 3. ∅ IR <sup>o</sup> 4. ∅ IR <sup>o</sup>	--
Ezpeleta 2023 <sup>24</sup>	n=80	12 weeks	1. ADF <sup>s</sup> 2. Ex <sup>j</sup> 3. ADF <sup>s</sup> + Ex <sup>g</sup> 4. Control	1. ↓5% 2. ↓2% 3. 5% ↓† 4. ∅	1. ↓ 2. ↓ 3. ↓† 4. ∅	1. ↓ 2. ↓ 3. ↓† 4. ∅	1. ↓ 2. ∅ 3. ↓† 4. ∅	1. ∅ 2. ∅ 3. ∅ SBP <sup>q</sup> ↓ DBP <sup>r</sup> 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ↓ 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ↓ 2. ∅ 3. ↓† 4. ∅	1. ↑ IS <sup>h</sup> ↓ IR <sup>o</sup> 2. ↑ IS <sup>h</sup> 3. ↑† IS <sup>h</sup> ↓ IR <sup>o</sup> 4. ∅ ∅ A1c <sup>a</sup>	--
Time Restricted Eating															
Moro 2020 <sup>27</sup>	n=16	4 weeks	1. TRE <sup>t</sup> 2. ND <sup>u</sup>	1. 2% ↓† 2. ∅	1. ↓ <sup>^</sup> 2. ∅	1. ∅ 2. ∅	--	--	--	--	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	--	1. ∅ differences between performance tests, ↑ peak power/BW <sup>v</sup> 2. ∅
Richardson 2023 <sup>28</sup>	n=24 n=15 analyzed	4 weeks	1. TRE <sup>t</sup> 2. ND <sup>u</sup>	1. ∅ 2. ∅	1. ↓ 2. ∅	1. ∅ 2. ∅	--	1. ∅ 2. ∅ SBP <sup>q</sup> / DBP <sup>r</sup>	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ IR <sup>o</sup> , IS 2. ∅ IR <sup>o</sup> , IS <sup>h</sup>	--
Correia 2023 <sup>29</sup>	n= 18	4 weeks	1. TRE <sup>t</sup> + RT <sup>w</sup> 2. ND <sup>u</sup> + RT <sup>w</sup>	1. ∅ 2. ∅	1. ↓ 2. ↓	1. ∅ 2. ∅	--	--	--	--	--	--	--	--	1. ↑ explosive upper body strength, ↑† bench press throw peak force & bench

															press throw dynamic index 2. ↑ explosive upper body strength, ↑ <sup>†</sup> lower and upper body muscle strength
Waldman 2023 <sup>30</sup>	n=15 n=12 analyzed	4 weeks	1. TRE <sup>t</sup>	1. ↓3%	1. ↓	1. ∅	--	--	1. ∅	1. ↑	1. ∅	1. ↓	1. ∅	∅	∅
Haganes 2022 <sup>33</sup>	n=131	7 weeks	1. TRE <sup>t</sup> 2. HIIT <sup>x</sup> 3. TRE <sup>t</sup> and HIIT <sup>x</sup> 4. Control	1. ↓2% <sup>†</sup> 2. ↓2% <sup>†</sup> 3. ↓4% <sup>†</sup> 4. ∅	1. ↓ <sup>†</sup> 2. ↓ <sup>†</sup> 3. ↓ <sup>†</sup> 4. ∅	1. ∅ ↓ <sup>†</sup> 2. ∅ 3. ∅ 4. ∅	1. ↓ <sup>†</sup> 2. ↓ <sup>†</sup> 3. ↓ <sup>†</sup> 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ↓ <sup>†</sup> 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ 2. ∅ 3. ∅ 4. ∅	1. ∅ A1c <sup>a</sup> , IR <sup>o</sup> 2. ∅ A1c <sup>a</sup> , IR <sup>o</sup> 3. ↓ A1c <sup>a</sup> , ∅IR <sup>o</sup> 4. ∅ A1c <sup>a</sup> , IR <sup>o</sup>	1. ∅ 2. ↑ V02peak <sup>l</sup> 3. ↑ V02peak <sup>l</sup> 4. ∅	
Moro 2016 <sup>31</sup>	n=34	8 weeks	1. TRE <sup>t</sup> + RT <sup>w</sup> 2. ND <sup>u</sup> + RT <sup>w</sup>	1. ∅ 2. ∅	1. ↓ <sup>†</sup> 2. ∅	1. ∅ 2. ∅	--	--	1. ∅ 2. ∅	1. ↑ 2. ∅	1. ↓ <sup>†</sup> 2. ∅	1. ↓ 2. ∅	1. ↓ 2. ∅	1. ↓ IR <sup>o</sup> 2. ∅	1. ↑ Leg press, hip sled 2. ↑ Leg press, hip sled
Moro 2021 <sup>42</sup>	n=20	52-week follow-up	1. TRE <sup>t</sup> 2. ND <sup>u</sup>	1. ↓3% <sup>†</sup> 2. ↑3%	1. ↓ <sup>†</sup> 2. ∅	1. ∅ 2. ∅ ↑ <sup>†</sup>	1. ∅ 2. ∅		1. ↓ 2. ∅	1. ↑ <sup>†</sup> 2. ∅	1. ↓ <sup>†</sup> 2. ∅	1. ↓ <sup>†</sup> 2. ∅	1. ↓ <sup>†</sup> 2. ∅	1. ↓ <sup>†</sup> IR <sup>o</sup> 2. ∅ IR <sup>o</sup>	1. ↓ <sup>†</sup> Thigh and arm circumference, ↑Bench and leg press 2. ↑Bench and leg press
Tinsley 2017 <sup>34</sup>	n=18	8 weeks	1. TRE <sup>t</sup> + RT <sup>w</sup> 2. ND <sup>u</sup> + RT <sup>w</sup>	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	--	--	--	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ IR <sup>o</sup> 2. ∅ IR <sup>o</sup>	1. ↑ Bench press, hip sled, hip sled endurance 2. ↑ Bench press, hip sled, hip sled endurance
Brady 2021 <sup>36</sup>	n=23 n=17 analyzed	8 weeks	1. TRE <sup>t</sup> 2. Control	1. ↓2% <sup>†</sup> 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	--	--	--	--	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ 2. ∅	1. ∅ IR <sup>o</sup> 2. ∅ IR <sup>o</sup>	1. ∅ V02peak <sup>l</sup> , FBLC <sup>y</sup> , HR at FBLC <sup>y</sup> , %HRmax <sup>k</sup> 2. ∅
Tinsley 2019 <sup>35</sup>	n=40	8 weeks	1. TRE <sup>t</sup> + RT <sup>w</sup> 2. TRE <sup>t</sup> + supplement + RT <sup>w</sup> 3. Control Diet + RT <sup>w</sup>	1. 1% <sup>†</sup> 2. 1% <sup>†</sup> 3. 2% <sup>†</sup>	1. ↓ 2. ↓ 3. ↓	1. ↑ 2. ↑ 3. ↑	--	1. ↓ DBP <sup>r</sup> 2. ↓ DBP <sup>r</sup> 3. ↓ DBP <sup>r</sup>	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	1. ∅ 2. ∅ 3. ∅	--	1. ↑ Max <sup>2</sup> strength and muscular performance 2. ↑ Max <sup>2</sup> strength and muscular performance 3. ↑ Max <sup>2</sup> strength and muscular performance



Kotarsky 2021 <sup>37</sup>	n=28 n=21 analyzed	8 weeks	1. TRE <sup>t</sup> 2. Control	1. ↓4% <sup>†</sup> 2. ∅	1. ↓ <sup>†</sup> 2. ∅	1. ↑ 2. ↑	1. ↓ 2. ↓ WC <sup>p</sup>	1. ∅ 2. ∅	--	1. ∅ 2. ∅	--	--	1. ∅ 2. ∅	1. ∅ 2. ∅ A1c <sup>a</sup>	1. ↑ knee flexion strength peak torque and endurance total work, dorsiflexion strength peak torque and endurance total work 2. ↑ knee flexion strength peak torque and endurance total work, dorsiflexion strength peak torque and endurance total work
Isemann 2021 <sup>38</sup>	n=35	8 weeks	1. TRE <sup>t</sup> 2. MBD <sup>aa</sup>	1. ↓5% 2. ↓5%	1. ↓ 2. ↓	1. ∅ 2. ∅	1. ↓ 2. ↓ WC <sup>p</sup>	--	--	--	--	--	--	--	--
Martínez- Rodríguez 2021 <sup>39</sup>	n=14	16 weeks	1. TRE <sup>t</sup> + HIIT <sup>x</sup> 2. HIIT <sup>x</sup>	1. ∅ 2. ∅	1. ↓ <sup>†</sup> 2. ∅	1. ∅ 2. ∅	--	--	--	--	--	--	--	--	1. ↑ counter- movement jump, relative mean power <sup>†</sup> 2. ∅

<sup>^</sup> P < 0.05, Significantly different between groups (between group effect).

<sup>†</sup> P < 0.05, Significant time by diet interaction.

<sup>a</sup>FM: Fat mass, <sup>b</sup>FFM: Fat free mass, <sup>c</sup>VF: Visceral fat mass, <sup>d</sup>LDL: Low density lipoprotein cholesterol, <sup>e</sup>HDL: High density lipoprotein cholesterol, <sup>f</sup>TG: Triglycerides, <sup>g</sup>IR: Insulin resistance, <sup>h</sup>IS: insulin sensitivity, <sup>i</sup>A1c: hemoglobin, <sup>j</sup>Ex: exercise, <sup>k</sup>HRmax: heart rate maximum, <sup>l</sup>V02peak: volume of oxygen peak, <sup>m</sup>CR: Calorie restriction, <sup>n</sup>Ad lib: Ad libitum energy intake, <sup>o</sup>SIT: sprint interval training, <sup>p</sup>WC: waist circumference, <sup>q</sup>SBP: Systolic blood pressure, <sup>r</sup>DBP: Diastolic blood pressure, <sup>s</sup>ADF: Alternate day fasting, <sup>t</sup>TRE: Time restricted eating, <sup>u</sup>ND: normal diet, <sup>v</sup>BW: body weight, <sup>w</sup>RT: Resistance training, <sup>x</sup>HIIT: high intensity interval training, <sup>y</sup>FBLC:fixed blood lactate concentration, <sup>z</sup>Max: maximum, <sup>aa</sup>MBD: macronutrient-based diet

**Table 3. Inclusion criteria (including sex, BMI, age, and training status), exercise modality, and adherence monitoring for the review of trials combining intermittent fasting with exercise.**

	Sex		BMI <sup>a</sup>			Age			Training status		Exercise			Adherence	
	M <sup>b</sup>	F <sup>c</sup>	<25	25-29.9	>30	18-45	45-65	>65	Trained	Untrained	AT <sup>d</sup>	RT <sup>e</sup>	HIIT <sup>f</sup>	Supervised	Wearable
<b>5:2</b>															
Batitucci 2022 <sup>16</sup>		●			●	●			--	--			●	●	
Keenan 2021, 2022 <sup>15,22</sup>	●	●	●	●	●	●	●	●		●		●		2 d/wk	
Kang 2022 <sup>17</sup>	●	●		●	●	●	●	●	--	--	●				
Hottenrott 2020 <sup>18</sup>	●	●		●	●	●	●		●		●				
Cooke 2022 <sup>21</sup>	●	●		●	●	●				●	●	●		●	
Headland 2019 <sup>19</sup>	●	●		●	●	●	●	●	--	--	●		●		
Jospe 2020 <sup>20</sup>	●	●		●	●	●	●	●	--	--	●	●	●		●
<b>ADF<sup>g</sup></b>															
Cho 2019 <sup>25,26</sup>	●	●		●	●	●	●		--	--	●	●		first week	
Bhutani 2013 <sup>23</sup>	●	●			●	●	●	●		●	●			●	
Ezpeleta 2023 <sup>24</sup>	●	●			●	●	●	●		●	●			●	
<b>TRE<sup>h</sup></b>															
Moro 2020 <sup>27</sup>	●		●			●			●		●				
Richardson 2023 <sup>28</sup>	●		●			●			●		●				
Correia 2023 <sup>29</sup>	●		●			●			●		●			1 d/wk	
Waldman 2023 <sup>30,43</sup>	●		●				●	●	●		●				
Haganes 2022 <sup>33</sup>	●	●		●	●	●				●			●	●	●
Moro 2016 <sup>31</sup>	●		●						●			●		●	
Moro 2021 <sup>42</sup>	●		●						●			●			
Tinsley 2017 <sup>34</sup>	●		●						●			●			
Brady 2021 <sup>36</sup>	●		●			●			●		●				●
Tinsley 2019 <sup>35</sup>		●	●			●			●			●		●	●
Kotarsky 2021 <sup>37</sup>	●	●		●	●	●	●			●	●	●		●	●
Isenmann 2021 <sup>38</sup>	●	●		●	●	●			●		●	●			
Martínez-Rodríguez 2021 <sup>39</sup>		●	●			●			●				●	●	

<sup>a</sup>BMI: Body Mass Index, <sup>b</sup>M: Male, <sup>c</sup>F: Female, <sup>d</sup>AT: Aerobic training, <sup>e</sup>RT: Resistance training, <sup>f</sup>HIIT: High intensity interval training, <sup>g</sup>ADF: Alternate day Fasting, <sup>h</sup>TRE: Time restricted eating